Not every baby gets all these symptoms and they can appear in any order.



Meningitis Baby Watch

Is your baby getting worse fast? Babies can get ill very quickly, so check often.

For information and support call our free helpline

080 8800 3344

helpline@meningitis.org www.meningitis.org



Meningitis & Septicaemia

strike fast and can kill

What should I do if I am worried?

Trust your instincts. If you think your baby has meningitis or septicaemia get medical help immediately. Say that you are worried it could be meningitis or septicaemia.

Know the symptoms

Early symptoms of meningitis and septicaemia (the blood poisoning form of the disease) can be like other childhood illnesses, but a baby will usually get ill quickly and get worse fast.

Watch out for tiny red or brown pin prick marks which can change into purple blotches or blood blisters. If your baby gets a rash, do the 'Tumbler Test'.

This rash can be harder to see on dark skin, so check for spots over baby's whole body, especially on paler areas like palms of the hands, the soles of the feet, on the tummy, inside the eyelids, and on the roof of the mouth.

How can I protect my baby?

Get your baby vaccinated. Vaccines give excellent protection but can't prevent all forms, so know the symptoms.

Tumbler test

If a glass tumbler is pressed firmly against a septicaemic rash, the marks will not fade. You will be able to see them through the glass. If this happens get medical help immediately.

Remember, a very ill baby needs medical help even if they have only a few spots, a rash that fades or no rash at all







For information and support call our free helpline.

UK **080 8800 3344** Ireland **1800 41 33 44**

Bristol 0333 405 6262 info@meningitis.org

Edinburgh 0131 510 2345 edinburgh@meningitis.org

Belfast 028 9032 1283 belfast@meningitis.org

Dublin 01 819 6931 dublin@meningitis.org

